

University of Groningen

Evaluation of π - π interactions in proteins using Trp analogs

Shao, Jinfeng

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Shao, J. (2016). *Evaluation of π - π interactions in proteins using Trp analogs: From protein labeling to quantitating the energies involved*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

- Abbott, G. L., Blouse, G. E., Perron, M. J., Shore, J. D., Luck, L. A. & Szabo, A. G. (2004). F-19 NMR studies of plasminogen activator inhibitor-1. *Biochemistry* **43**, 1507-1519.
- Afonine, P. V., Grosse-Kunstleve, R. W., Echols, N., Headd, J. J., Moriarty, N. W., Mustyakimov, M., Terwilliger, T. C., Urzhumtsev, A., Zwart, P. H. & Adams, P. D. (2012). Towards automated crystallographic structure refinement with phenix.refine. *Acta Crystallogr D* **68**, 352-367.
- Agustiandari, H., Lubelski, J., van den Berg van Saparoea, H. B., Kuipers, O. P. & Driessen, A. J. (2008). LmrR is a transcriptional repressor of expression of the multidrug ABC transporter LmrCD in *Lactococcus lactis*. *J Bacteriol* **190**, 759-763.
- Aller, K., Adamberg, K., Timarova, V., Seiman, A., Festsenko, D. & Vilu, R. (2014). Nutritional requirements and media development for *Lactococcus lactis* IL1403. *Appl Microbiol Biotechnol* **98**, 5871-5881.
- Anslyn, E. V. & Dougherty, D. A. (2006). *Modern physical organic chemistry*, pp. 180-185. Sausalito, CA: University Science.
- Antonczak, A. K., Simova, Z., Yonemoto, I. T., Bochtler, M., Piasecka, A., Czapinska, H., Brancale, A. & Tippmann, E. M. (2011). Importance of single molecular determinants in the fidelity of expanded genetic codes. *P Natl Acad Sci USA* **108**, 1320-1325.
- Asensio, J. L., Ardá, A., Cañada, F. J. & Jiménez-Barbero, J. (2013). Carbohydrate-aromatic interactions. *Acc Chem Res* **46**, 946-954.
- Askew, B., Ballester, P., Buhr, C., Jeong, K. S., Jones, S., Parris, K., Williams, K. & Rebek, J. (1989). Molecular Recognition with Convergent Functional-

Groups .6. Synthetic and Structural Studies with a Model Receptor for Nucleic-Acid Components. *J Am Chem Soc* **111**, 1082-1090.

Aziel, R. & Gazit, E. (2001). Analysis of the minimal amyloid-forming fragment of the islet amyloid polypeptide: an experimental support for the key role of the phenylalanine residue in amyloid formation. *J Biol Chem* **276**, 34156-34161.

Balbach, J. J., Ishii, Y., Antzutkin, O. N., Leapman, R. D., Rizzo, N. W., Dyda, F., Reed, J. & Tycko, R. (2000). Amyloid fibril formation by A β 16-22 , a seven-residue fragment of the alzheimer's β -amyloid peptide, and structural characterization by solid state NMR. *Biochemistry* **39**, 13748-13759.

Barnard, E. A., Miledi, R. & Sumikawa, K. (1982). Translation of exogenous messenger-RNA coding for nicotinic acetylcholine-receptors produces functional receptors in *Xenopus* oocytes. *Proc R Soc Ser B-Bio* **215**, 241-246.

Barone, G., Guerra, C. F., Gambino, N., Silvestri, A., Lauria, A., Almerico, A. M. & Bickelhaupt, F. M. (2008). Intercalation of daunomycin into stacked DNA base pairs. DFT study of an anticancer drug. *J Biomol Struct Dyn* **26**, 115-129.

Barthwal, R., Sharma, U., Srivastava, N., Jain, M., Awasthi, P., Kaur, M., Barthwal, S. & Govil, G. (2006). Structure of daunomycin complexed to d-TGATCA by two-dimensional nuclear magnetic resonance spectroscopy. *Eur J Med Chem* **41**, 27-39.

Berntsson, R. P., Alia Oktaviani, N., Fusetti, F., Thunnissen, A. M., Poolman, B. & Slotboom, D. J. (2009). Selenomethionine incorporation in proteins expressed in *Lactococcus lactis*. *Protein Sci* **18**, 1121-1127.

Brejc, K., van Dijk, W. J., Klaassen, R. V., Schuurmans, M., van der Oost, J., Smit, A. B. & Sixma, T. K. (2001). Crystal structure of an ACh-binding protein

- reveals the ligand-binding domain of nicotinic receptors. *Nature* **411**, 269-276.
- Broos, J., ter Veld, F. & Robillard, G. T. (1999). Membrane protein-ligand interactions in *Escherichia coli* vesicles and living cells monitored *via* a biosynthetically incorporated tryptophan analogue. *Biochemistry* **38**, 9798-9803.
- Broos, J., Maddalena, F. & Hesp, B. H. (2004). *In vivo* synthesized proteins with monoexponential fluorescence decay kinetics. *J Am Chem Soc* **126**, 22-23.
- Broos, J. (2014). Biosynthetic incorporation of tryptophan analogs in proteins. *Methods Mol Biol* **1076**, 359-370.
- Budisa, N. (2004). Prolegomena to future experimental efforts on genetic code engineering by expanding its amino acid repertoire. *Angew Chem Int Ed* **43**, 6426-6463.
- Burley, S. & Petsko, G. (1985). Aromatic-aromatic interaction: a mechanism of protein structure stabilization. *Science* **229**, 23-28.
- Butterfield, S. M., Patel, P. R. & Waters, M. L. (2002). Contribution of aromatic interactions to α -helix stability. *J Am Chem Soc* **124**, 9751-9755.
- Callis, P. R. (1997). L-1(a) and L-1(b) transitions of tryptophan: Applications of theory and experimental observations to fluorescence of proteins. *Methods Enzymol* **278**, 113-150.
- Carver, F. J., Hunter, C. A., Jones, P. S., Livingstone, D. J., McCabe, J. F., Seward, E. M., Tiger, P. & Spey, S. E. (2001). Quantitative measurements of edge-to-face aromatic interactions by using chemical double-mutant cycles. *Chem Eur J* **7**, 4854-4862.

- Carver, F. J., Hunter, C. A., Livingstone, D. J., McCabe, J. F. & Seward, E. M. (2002). Substituent effects on edge-to-face aromatic interactions. *Chem Eur J* **8**, 2847-2859.
- Černý, J. i., Kabeláč, M. & Hobza, P. (2008). Double-helical → ladder structural transition in the b-DNA is induced by a loss of dispersion energy. *J Am Chem Soc* **130**, 16055-16059.
- Chaires, J. B., Dattagupta, N. & Crothers, D. M. (1985). Kinetics of the daunomycin-DNA interaction. *Biochemistry* **24**, 260-267.
- Chakravorty, D. K., Wang, B., Ucisik, M. N. & Merz, K. M. (2011). Insight into the cation- π interaction at the metal binding site of the copper metallochaperone CusF. *J Am Chem Soc* **133**, 19330-19333.
- Chen, V. B., Arendall, W. B., Headd, J. J., Keedy, D. A., Immormino, R. M., Kapral, G. J., Murray, L. W., Richardson, J. S. & Richardson, D. C. (2010). MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallogr D* **66**, 12-21.
- Ciatto, C., D'Amico, M. L., Natile, G., Secco, F. & Venturini, M. (1999). Intercalation of proflavine and a platinum derivative of proflavine into double-helical poly(A). *Biophysical J* **77**, 2717-2724.
- Connor, R. E. & Tirrell, D. A. (2007). Non-canonical amino acids in protein polymer design. *Polym Rev* **47**, 9-28.
- Cooke, G. & Rotello, V. M. (2002). Methods of modulating hydrogen bonded interactions in synthetic host-guest systems. *Chem Soc Rev* **31**, 275-286.
- Cox, E. G., Cruickshank, D. W. J. & Smith, J. A. S. (1958). The Crystal Structure of Benzene at -3-Degrees-C. *Proc R Soc Lon Ser-A* **247**, 1-21.

- Cozzi, F. & Siegel, J. S. (1995). Interaction between stacked aryl groups in 1,8-diarylnaphthalenes: Dominance of polar/ π over charge-transfer effects. *Pure Appl. Chem* **67**, 683-689.
- Cubberley, M. S. & Iverson, B. L. (2001). ^1H NMR Investigation of Solvent Effects in Aromatic Stacking Interactions. *J Am Chem Soc* **123**, 7560-7563.
- D'Amico, M. L., Paiotta, V., Secco, F. & Venturini, M. (2002). A kinetic study of the intercalation of ethidium bromide into poly(A)·poly(U). *J Phys Chem B* **106**, 12635-12641.
- Daeffler, K. N. M., Lester, H. A. & Dougherty, D. A. (2012). Functionally important aromatic-aromatic and sulfur- π interactions in the D2 dopamine receptor. *J Am Chem Soc* **134**, 14890-14896.
- Dalvi, V. H. & Rosicky, P. J. (2010). Molecular origins of fluorocarbon hydrophobicity. *P Natl Acad Sci USA* **107**, 13603-13607.
- Davies, D. B., Eaton, R. J., Baranovsky, S. F. & Veselkov, A. N. (2000). NMR investigation of the complexation of daunomycin with deoxytetranucleotides of different base sequence in aqueous solution. *J Biomol Struct Dyn* **17**, 887-901.
- Del Rizzo, P. A., Krishnan, S. & Trievel, R. C. (2012). Crystal structure and functional analysis of JMJD5 indicate an alternate specificity and function. *Mol Cell Biol* **32**, 4044-4052.
- Doublet, S., Bricogne, G., Gilmore, C. & Carter, C. W. (1995). Tryptophanyl-transfer-RNA synthetase crystal-structure reveals an unexpected homology to tyrosyl-transfer-RNA synthetase. *Structure* **3**, 17-31.
- Dougherty, D. A. (2013). The cation- π interaction. *Acc Chem Res* **46**, 885-893.

- Dougherty, D. A. & Van Arnem, E. B. (2014). *In vivo* incorporation of non-canonical amino acids by using the chemical aminoacylation strategy: a broadly applicable mechanistic tool. *Chembiochem* **15**, 1710-1720.
- El Khattabi, M., van Roosmalen, M. L., Jager, D., Metselaar, H., Permentier, H., Leenhouts, K. & Broos, J. (2008). *Lactococcus lactis* as expression host for the biosynthetic incorporation of tryptophan analogues into recombinant proteins. *Biochem J* **409**, 193-198.
- Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. (2010). Features and development of Coot. *Acta Crystallogr D* **66**, 486-501.
- Evans, P. R. & Murshudov, G. N. (2013). How good are my data and what is the resolution? *Acta Crystallogr D* **69**, 1204-1214.
- Fersht, A. R. (1999). *Structure and mechanism in protein science : a guide to enzyme catalysis and protein folding*, pp. 324-347. New York, N.Y. : Freeman.
- Fishman, R., Ankilova, V., Moor, N. & Safro, M. (2001). Structure at 2.6 angstrom resolution of phenylalanyl-tRNA synthetase complexed with phenylalanyl-adenylate in the presence of manganese. *Acta Crystallogr Sect D* **57**, 1534-1544.
- Frenkel-Pinter, M., Tal, S., Scherzer-Attali, R., Abu-Hussien, M., Alyagor, I., Eisenbaum, T., Gazit, E. & Segal, D. (2016). Naphthoquinone-Tryptophan Hybrid Inhibits Aggregation of the Tau-Derived Peptide PHF6 and Reduces Neurotoxicity. *J Alzheimers Dis* **51**, 165-178.
- Friedman, R. A. & Honig, B. (1996). New evidence that the hydrophobic effect and dispersion are not major driving forces for nucleotide base stacking- Response. *Biophys J* **71**, 3525-3526.

- Gazit, E. (2002). A possible role for pi-stacking in the self-assembly of amyloid fibrils. *Faseb J* **16**, 77-83.
- Gellman, S. H., Haque, T. S. & Newcomb, L. F. (1996). New evidence that the hydrophobic effect and dispersion are not major driving forces for nucleotide base stacking. *Biophys J* **71**, 3523-3525.
- Geng, Y., Takatani, T., Hohenstein, E. G. & Sherrill, C. D. (2010). Accurately characterizing the π - π interaction energies of indole-benzene complexes. *J Phys Chem A* **114**, 3576-3582.
- González-Menéndez, P., García-Ocaña, M. & de los Toyos, J. R. (2013). A deeper analysis of the epitope/paratope of PLY-5, a mouse monoclonal antibody which recognises the conserved undecapeptide tryptophan-rich loop (ECTGLAWWWWR) of bacterial cholesterol-dependent cytolysins. *Biochem Biophys Res Commun* **430**, 14-19.
- Goodman, A. J., Breinlinger, E. C., McIntosh, C. M., Grimaldi, L. N. & Rotello, V. M. (2001). Model systems for flavoenzyme activity. Control of flavin recognition via specific electrostatic interactions. *Org Lett* **3**, 1531-1534.
- Grimme, S. (2008). Do special noncovalent π - π stacking interactions really exist? *Angew Chem Int Ed* **47**, 3430-3434.
- Grishin, N. V., Osterman, A. L., Brooks, H. B., Phillips, M. A. & Goldsmith, E. J. (1999). X-ray structure of ornithine decarboxylase from *Trypanosoma brucei*: the native structure and the structure in complex with α -difluoromethylornithine. *Biochemistry* **38**, 15174-15184.
- Guckian, K. M., Schweitzer, B. A., Ren, R. X. F., Sheils, C. J., Tahmassebi, D. C. & Kool, E. T. (2000). Factors contributing to aromatic stacking in water: evaluation in the context of DNA. *J Am Chem Soc* **122**, 2213-2222.

- Guo, R. T. (2003). Crystal structure of octaprenyl pyrophosphate synthase from hyperthermophilic *thermotoga maritima* and mechanism of product chain length determination. *J Biol Chem* **279**, 4903-4912.
- Gurdon, J. B., Lane, C. D., Woodland, H. R. & Marbaix, G. (1971). Use of frog eggs and oocytes for study of messenger RNA and its translation in living cells. *Nature* **233**, 177-182.
- Hill, R. B. & DeGrado, W. F. (1998). Solution structure of α 2D, a nativelike *de novo* designed protein. *J Am Chem Soc* **120**, 1138-1145.
- Hogue, C. W., Rasquinha, I., Szabo, A. G. & MacManus, J. P. (1992). A new intrinsic fluorescent probe for proteins. Biosynthetic incorporation of 5-hydroxytryptophan into oncomodulin. *Febs Lett* **310**, 269-272.
- Huang, R., Wu, L., McElheny, D., Bouř, P., Roy, A. & Keiderling, T. A. (2009). Cross-strand coupling and site-specific unfolding thermodynamics of a trpzip β -hairpin peptide using ^{13}C isotopic labeling and IR spectroscopy. *J Phys Chem B* **113**, 5661-5674.
- Hudson, B. S., Harris, D. L., Ludescher, R. D., Cooney-Freed, A. & Cavalier, S. (1986). *Applications of fluorescence in the biomedical sciences*, edited by D. L. Taylor & L. Abbott, pp. 159-202. New York: A.R. Liss.
- Hudson, K. L., Bartlett, G. J., Diehl, R. C., Agirre, J., Gallagher, T., Kiessling, L. L. & Woolfson, D. N. (2015). Carbohydrate–aromatic interactions in proteins. *J Am Chem Soc* **137**, 15152-15160.
- Hunter, C. A. & Sanders, J. K. M. (1990). The nature of pi-pi interactions. *J Am Chem Soc* **112**, 5525-5534.
- Hunter, C. A., Lawson, K. R., Perkins, J. & Urch, C. J. (2001). Aromatic interactions. *J Chem Soc Perk T 2*, 651-669.

- Jensen, P. R. & Hammer, K. (1993). Minimal requirements for exponential-growth of *Lactococcus-Lactis*. *Appl Environ Microbiol* **59**, 4363-4366.
- Kabsch, W. (2010). Xds. *Acta Crystallogr D* **66**, 125-132.
- Kim, E., Paliwal, S. & Wilcox, C. S. (1998). Measurements of molecular electrostatic field effects in edge-to-face aromatic interactions and CH- π interactions with implications for protein folding and molecular recognition. *J Am Chem Soc* **120**, 11192-11193.
- Klosterman, J. K., Yamauchi, Y. & Fujita, M. (2009). Engineering discrete stacks of aromatic molecules. *Chem Soc Rev* **38**, 1714.
- Kryger, G., Silman, I. & Sussman, J. L. (1999). Structure of acetylcholinesterase complexed with E2020 (Aricept): implications for the design of new anti-Alzheimer drugs. *Structure* **7**, 297-307.
- Kuipers, O. P., Beerthuyzen, M. M., Deruyter, P. G. G. A., Luesink, E. J. & Devos, W. M. (1995). Autoregulation of nisin biosynthesis in *Lactococcus-Lactis* by signal-transduction. *J Biol Chem* **270**, 27299-27304.
- Kuipers, O. P., de Ruyter, P. G., Kleerebezem, M. & de Vos, W. M. (1997). Controlled overproduction of proteins by lactic acid bacteria. *Trends Biotechnol* **15**, 135-140.
- Kunji, E. R. S., Slotboom, D. J. & Poolman, B. (2003). *Lactococcus lactis* as host for overproduction of functional membrane proteins. *Biochim Biophys Acta-Biomem* **1610**, 97-108.
- Kwon, I., Wang, P. & Tirrell, D. A. (2006). Design of a bacterial host for site-specific incorporation of p-bromophenylalanine into recombinant proteins. *J Am Chem Soc* **128**, 11778-11783.

- Kwon, I. & Tirrell, D. A. (2007). Site-specific incorporation of tryptophan analogues into recombinant proteins in bacterial cells. *J Am Chem Soc* **129**, 10431-10437.
- Langner, K. M., Kedzierski, P., Sokalski, W. A. & Leszczynski, J. (2006). Physical nature of ethidium and proflavine interactions with nucleic acid bases in the intercalation plane. *J Phys Chem B* **110**, 9720-9727.
- Lashuel, H. A., Hartley, D. M., Balakhaneh, D., Aggarwal, A., Teichberg, S. & Callaway, D. J. E. (2002). New class of inhibitors of amyloid-beta fibril formation - Implications for the mechanism of pathogenesis in Alzheimer's disease. *J Biol Chem* **277**, 42881-42890.
- Laughrey, Z. R., Kiehna, S. E., Riemen, A. J. & Waters, M. L. (2008). Carbohydrate- π interactions: what are they worth? *J Am Chem Soc* **130**, 14625-14633.
- Lepthien, S., Hoesl, M. G., Merkel, L. & Budisa, N. (2008). Azatryptophans endow proteins with intrinsic blue fluorescence. *P Natl Acad Sci USA* **105**, 16095-16100.
- Li, H. J. & Crothers, D. M. (1969). Relaxation studies of the proflavine-DNA complex: the kinetics of an intercalation reaction. *J Mol Biol* **39**, 461-477.
- Li, S., Cooper, V. R., Thonhauser, T., Lundqvist, B. I. & Langreth, D. C. (2009). Stacking interactions and DNA intercalation. *J Phys Chem B* **113**, 11166-11172.
- Liu, T. Q., Callis, P. R., Hesp, B. H., de Groot, M., Buma, W. J. & Broos, J. (2005). Ionization potentials of fluoroindoles and the origin of nonexponential tryptophan fluorescence decay in proteins. *J Am Chem Soc* **127**, 4104-4113.

- Liyanage, W. & Nilsson, B. L. (2016). Substituent Effects on the Self-Assembly/Coassembly and Hydrogelation of Phenylalanine Derivatives. *Langmuir* **32**, 787-799.
- Loidl, G., Musiol, H. J., Budisa, N., Huber, R., Poirot, S., Fourmy, D. & Moroder, L. (2000). Synthesis of beta-(1-amulenyl)-L-alanine as a potential blue-colored fluorescent tryptophan analog and its use in peptide synthesis. *J Pept Sci* **6**, 139-144.
- Ma, J. C. & Dougherty, D. A. (1997). The cation- π interaction. *Chem Rev* **97**, 1303-1324.
- Madoori, P. K., Agustiandari, H., Driessen, A. J. M. & Thunnissen, A. (2009). Structure of the transcriptional regulator LmrR and its mechanism of multidrug recognition. *EMBO J* **28**, 156-166.
- Makin, O. S., Atkins, E., Sikorski, P., Johansson, J. & Serpell, L. C. (2005). Molecular basis for amyloid fibril formation and stability. *P Natl Acad Sci USA* **102**, 315-320.
- Martinez, C. R. & Iverson, B. L. (2012). Rethinking the term "pi-stacking". *Chem Sci* **3**, 2191-2201.
- McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C. & Read, R. J. (2007). Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674.
- Mirassou, Y., Santiveri, C. M., Pérez de Vega, M. J., González-Muñiz, R. & Jiménez, M. A. (2009). Disulfide bonds versus trp...trp pairs in irregular β -hairpins: NMR structure of vamin loop 3-derived peptides as a case study. *ChemBioChem* **10**, 902-910.

- Morello, E., Bermudez-Humaran, L. G., Llull, D., Sole, V., Miraglio, N., Langella, P. & Poquet, I. (2008). *Lactococcus lactis*, an efficient cell factory for recombinant protein production and secretion. *J Mol Microbiol Biotechnol* **14**, 48-58.
- Moroz, Y. S., Binder, W., Nygren, P., Caputo, G. A. & Korendovych, I. V. (2013). Painting proteins blue: beta-(1-azulenyl)-L-alanine as a probe for studying protein-protein interactions. *Chem Commun* **49**, 490-492.
- Muehldorf, A. V., Vanengen, D., Warner, J. C. & Hamilton, A. D. (1988). Aromatic Aromatic Interactions in Molecular Recognition - a Family of Artificial Receptors for Thymine That Shows Both Face-to-Face and Edge-to-Face Orientations. *J Am Chem Soc* **110**, 6561-6562.
- Mukherjee, A., Lavery, R., Bagchi, B. & Hynes, J. T. (2008). On the molecular mechanism of drug intercalation into DNA: A simulation study of the intercalation pathway, free energy, and DNA structural changes. *J Am Chem Soc* **130**, 9747-9755.
- Muller-Werkmeister, H. M. & Bredenbeck, J. (2014). A donor-acceptor pair for the real time study of vibrational energy transfer in proteins. *Phys Chem Chem Phys* **16**, 3261-3266.
- Newcomb, L. F. & Gellman, S. H. (1994). Aromatic stacking interactions in aqueous solution: evidence that neither classical hydrophobic effects nor dispersion forces are important. *J Am Chem Soc* **116**, 4993-4994.
- Nowak, M. W., Kearney, P. C., Sampson, J. R., Saks, M. E., Labarca, C. G., Silverman, S. K., Zhong, W., Thorson, J. S., Abelson, J. N., Davidson, N., Schultz, P. G., Dougherty, D. A. & Lester, H. A. (1995). Nicotinic receptor-binding site probed with unnatural amino-acid-incorporation in intact-cells. *Science* **268**, 439-442.
- Nowak, M. W., Gallivan, J. P., Silverman, S. K., Labarca, C. G., Dougherty, D. A. & Lester, H. A. (1998). *In vivo* incorporation of unnatural amino acids

- into ion channels in *Xenopus* oocyte expression system. *Methods Enzymol* **293**, 504-529.
- Okada, A., Miura, T. & Takeuchi, H. (2001). Protonation of histidine and histidine-tryptophan interaction in the activation of the M2 ion channel from influenza A virus. *Biochemistry* **40**, 6053-6060.
- Otto, R., Tenbrink, B., Veldkamp, H. & Konings, W. N. (1983). The relation between growth-rate and electrochemical proton gradient of *Streptococcus-Cremoris*. *Fems Microbiol Lett* **16**, 69-74.
- Pace, C. J., Zheng, H., Mylvaganam, R., Kim, D. & Gao, J. (2011). Stacked fluoroaromatics as supramolecular synthons for programming protein dimerization specificity. *Angew Chem Int Ed* **51**, 103-107.
- Pace, C. J. & Gao, J. (2013). Exploring and exploiting polar- π interactions with fluorinated aromatic amino acids. *Acc Chem Res* **46**, 907-915.
- Paliwal, S., Geib, S. & Wilcox, C. S. (1994). Chemistry of synthetic receptors and functional-group arrays .24. molecular torsion balance for weak molecular recognition forces - effects of tilted-t edge-to-face aromatic interactions on conformational selection and solid-state structure. *J Am Chem Soc* **116**, 4497-4498.
- Petrovic, D. M., Leenhouts, K., van Roosmalen, M. L., Kleinjan, F. & Broos, J. (2012). Monitoring lysin motif-ligand interactions via tryptophan analog fluorescence spectroscopy. *Anal Biochem* **428**, 111-118.
- Petrovic, D. M., Leenhouts, K., van Roosmalen, M. L. & Broos, J. (2013a). An expression system for the efficient incorporation of an expanded set of tryptophan analogues. *Amino Acids* **44**, 1329-1336.

- Petrovic, D. M., Hesp, B. H. & Broos, J. (2013b). Emitting state of 5-hydroxyindole, 5-hydroxytryptophan, and 5-hydroxytryptophan incorporated in proteins. *J Phys Chem B* **117**, 10792-10797.
- Poolman, B. & Konings, W. N. (1988). Relation of growth of *Streptococcus-Lactis* and *Streptococcus-Cremoris* to amino-acid transport. *J Bacteriol* **170**, 700-707.
- Porat, Y., Abramowitz, A. & Gazit, E. (2006). Inhibition of amyloid fibril formation by polyphenols: structural similarity and aromatic interactions as a common inhibition mechanism. *Chem Biol Drug Design* **67**, 27-37.
- Protozanova, E., Yakovchuk, P. & Frank-Kamenetskii, M. D. (2004). Stacked–unstacked equilibrium at the nick site of DNA. *J Mol Biol* **342**, 775-785.
- Quigley, G. J., Wang, A. H., Ughetto, G., van der Marel, G., van Boom, J. H. & Rich, A. (1980). Molecular structure of an anticancer drug-DNA complex: daunomycin plus d(CpGpTpApCpG). *P Natl Acad Sci USA* **77**, 7204-7208.
- Rahimi, M., van der Meer, J. Y., Geertsema, E. M., Poddar, H., Baas, B. J. & Poelarends, G. J. (2016). Mutations closer to the active site improve the promiscuous aldolase activity of 4-oxalocrotonate tautomerase more effectively than distant mutations. *Chembiochem* **17**, 1225-1228.
- Rashkin, M. J. & Waters, M. L. (2002). Unexpected substituent effects in offset π – π stacked interactions in water. *J Am Chem Soc* **124**, 1860-1861.
- Ross, J. B. A., Senear, D. F., Waxman, E., Kombo, B. B., Rusinova, E., Huang, Y. T., Laws, W. R. & Hasselbacher, C. A. (1992). Spectral enhancement of proteins-biological incorporation and fluorescence characterization of 5-hydroxytryptophan in bacteriophage-lambda ci repressor. *P Natl Acad Sci USA* **89**, 12023-12027.

- Ross, J. B. A., Szabo, A. G. & Hogue, C. W. (1997). Enhancement of protein spectra with tryptophan analogs: fluorescence spectroscopy of protein-protein and protein-nucleic acid interactions. *Methods Enzymol* **278**, 151-190.
- Ross, J. B. A., Rusinova, E., Luck, L. A. & Rousslang, K. W. (2000). *Topics in Fluorescence Spectroscopy: Volume 6: Protein Fluorescence*, edited by J. R. Lakowicz, pp. 17-42. Boston, MA: Springer US.
- Santiveri, C. M. & Jimenez, M. A. (2010). Tryptophan Residues: Scarce in Proteins but Strong Stabilizers of beta-Hairpin Peptides. *Biopolymers* **94**, 779-790.
- Scherzer-Attali, R., Shaltiel-Karyo, R., Adalist, Y. H., Segal, D. & Gazit, E. (2012). Generic inhibition of amyloidogenic proteins by two naphthoquinone-tryptophan hybrid molecules. *Proteins* **80**, 1962-1973.
- Schneider, H. J., Philippi, K. & Pohlmann, J. (1984). Host-Guest Chemistry .2. Structure-Dependence and Solvent-Dependence in the Complexation of Lipophilic Substrates in a Water-Soluble Azacyclophane. *Angew Chem Int Edit* **23**, 908-910.
- Shao, J., Korendovych, I. V. & Broos, J. (2015a). Biosynthetic incorporation of the azulene moiety in proteins with high efficiency. *Amino Acids* **47**, 213-216.
- Shao, J., Korendovych, I. V. & Broos, J. (2015b). Biosynthetic incorporation of the azulene moiety in proteins with high efficiency. *Amino Acids* **47**, 213-216.
- Shao, J., Marcondes, M. F., Oliveira, V. & Broos, J. (2016). Development of chemically defined media to express Trp-analog-labeled proteins in a *Lactococcus lactis* Trp auxotroph. *J Mol Microbiol Biotechnol* **26**, 269-276.

- Shi, X. Q. & Bartell, L. S. (1988). Electron-Diffraction and Monte-Carlo Studies of Liquids .1. Intermolecular Interactions for Benzene. *J Phys Chem-Us* **92**, 5667-5673.
- Shieh, H.-S., Berman, H. M., Dabrow, M. & Neidle, S. (1980). The structure of drug-deoxydinucleoside phosphate complex; generalized conformational behavior of intercalation complexes with RNA and DNA fragments. *Nucl Acids Res* **8**, 85-98.
- Soumillion, P., Jespers, L., Vervoort, J. & Fastrez, J. (1995). Biosynthetic incorporation of 7-azatryptophan into the phage lambda lysozyme: estimation of tryptophan accessibility, effect on enzymatic activity and protein stability. *Protein Eng* **8**, 451-456.
- Speight, L. C., Samanta, M. & Petersson, E. J. (2014). Minimalist approaches to protein labelling: getting the most fluorescent bang for your steric buck. *Aust J Chem* **67**, 686-700.
- Šponer, J., Riley, K. E. & Hobza, P. (2008). Nature and magnitude of aromatic stacking of nucleic acid bases. *Phys Chem Chem Phys* **10**, 2595.
- Šponer, J., Šponer, J. E., Mládek, A., Jurečka, P., Banáš, P. & Otyepka, M. (2013). Nature and magnitude of aromatic base stacking in DNA and RNA: quantum chemistry, molecular mechanics and experiment. *Biopolymers* **99**, 978-988.
- Subra, A. K., Nissen, M. S., Lewis, K. M., Muralidharan, A. K., Sanchez, E. J., Milting, H. & Kang, C. (2012). Molecular mechanisms of pharmaceutical drug binding into calsequestrin. *Int J Mol Sci* **13**, 14326-14343.
- Sumikawa, K., Houghton, M., Emtage, J. S., Richards, B. M. & Barnard, E. A. (1981). Active multi-subunit Ach receptor assembled by translation of heterologous messenger-RNA in *Xenopus* oocytes. *Nature* **292**, 862-864.

- Takekiyo, T., Wu, L., Yoshimura, Y., Shimizu, A. & Keiderling, T. A. (2009). Relationship between hydrophobic interactions and secondary structure stability for trpzip β -hairpin peptides *Biochemistry* **48**, 1543-1552.
- Takeuchi, K., Tokunaga, Y., Imai, M., Takahashi, H. & Shimada, I. (2014). Dynamic multidrug recognition by multidrug transcriptional repressor LmrR. *Sci Rep* **4**, 6922.
- Tenidis, K., Waldner, M., Bernhagen, J., Fischle, W., Bergmann, M., Weber, M., Merkle, M.-L., Voelter, W., Brunner, H. & Kapurniotu, A. (2000). Identification of a penta- and hexapeptide of islet amyloid polypeptide (IAPP) with amyloidogenic and cytotoxic properties. *J Mol Biol* **295**, 1055-1071.
- Tsuzuki, S., Honda, K., Uchamaru, T., Mikami, M. & Tanabe, K. (2002). Origin of attraction and directionality of the π/π interaction: model chemistry calculations of benzene dimer interaction. *J Am Chem Soc* **124**, 104-112.
- Turnell, W. G. & Finch, J. T. (1992). Binding of the dye congo red to the amyloid protein pig insulin reveals a novel homology amongst amyloid-forming peptide sequences. *J Mol Biol* **227**, 1205-1223.
- Twine, S. M. & Szabo, A. G. (2003). Fluorescent amino acid analogs. *Methods Enzymol* **360**, 104-127.
- Twine, S. M., Murphy, L., Phillips, R. S., Callis, P., Cash, M. T. & Szabo, A. G. (2003). The photophysical properties of 6-azaindole. *J Phys Chem B* **107**, 637-645.
- van der Berg, J. P., Madoori, P. K., Komarudin, A. G., Thunnissen, A. M. & Driessen, A. J. M. (2015). Binding of the Lactococcal drug dependent

transcriptional regulator LmrR to its ligands and responsive promoter regions. *Plos One* **10**, e0135467.

Wang, A. H. J., Ughetto, G., Quigley, G. J. & Rich, A. (1987). Interactions between an anthracycline antibiotic and DNA: molecular structure of daunomycin complexed to d(CpGpTpApCpG) at 1.2 Å resolution. *Biochemistry* **26**, 1152-1163.

Wang, H., Zhang, H., Zou, Y., Mi, Y., Lin, S., Xie, Z., Yan, Y. & Zhang, H. (2014). Structural insight into the tetramerization of an iterative ketoreductase SiaM through aromatic residues in the interfaces. *PLoS ONE* **9**, e97996.

Waters, M. L. (2002). Aromatic interactions in model systems. *Curr Opin Chem Biol* **6**, 736-741.

Wheeler, S. E. (2011). Local nature of substituent effects in stacking interactions. *J Am Chem Soc* **133**, 10262-10274.

Wheeler, S. E. (2013). Understanding substituent effects in noncovalent interactions involving aromatic rings. *Accounts Chem Res* **46**, 1029-1038.

Wheeler, S. E. (2015). *Noncovalent Forces*, pp. 421-442. Switzerland: Springer International Publishing.

Wilhelm, M., Mukherjee, A., Bouvier, B., Zakrzewska, K., Hynes, J. T. & Lavery, R. (2012). Multistep drug intercalation: molecular dynamics and free energy studies of the binding of daunomycin to DNA. *J Am Chem Soc* **134**, 8588-8596.

Winn, M. D., Ballard, C. C., Cowtan, K. D., Dodson, E. J., Emsley, P., Evans, P. R., Keegan, R. M., Krissinel, E. B., Leslie, A. G. W., McCoy, A., McNicholas, S. J., Murshudov, G. N., Pannu, N. S., Potterton, E. A., Powell, H. R., Read,

- R. J., Vagin, A. & Wilson, K. S. (2011). Overview of the CCP4 suite and current developments. *Acta Crystallogr D* **67**, 235-242.
- Wu, L., McElheny, D., Huang, R. & Keiderling, T. A. (2009). Role of tryptophan-tryptophan interactions in trpzip β -hairpin formation, structure, and stability. *Biochemistry* **48**, 10362-10371.
- Xie, J. M. & Schultz, P. G. (2006). Innovation: a chemical toolkit for proteins-an expanded genetic code. *Nat Rev Mol Cell Bio* **7**, 775-782.
- Xing, L., Kurumbail, R. G., Frazier, R. B., Davies, M. S., Fujiwara, H., Weinberg, R. A., Gierse, J. K., Caspers, N., Carter, J. S., McDonald, J. J., Moore, W. M. & Vazquez, M. L. (2008). Homo-timeric structural model of human microsomal prostaglandin E synthase-1 and characterization of its substrate/inhibitor binding interactions. *J Comput Aid Mol Des* **23**, 13-24.
- Xiu, X. A., Puskar, N. L., Shanata, J. A. P., Lester, H. A. & Dougherty, D. A. (2009). Nicotine binding to brain receptors requires a strong cation- π interaction. *Nature* **458**, 534-537.
- Yakovchuk, P. (2006). Base-stacking and base-pairing contributions into thermal stability of the DNA double helix. *Nucleic Acids Res* **34**, 564-574.
- Yang, F. S., Lim, G. P., Begum, A. N., Ubeda, O. J., Simmons, M. R., Ambegaokar, S. S., Chen, P. P., Kaye, R., Glabe, C. G., Frautschy, S. A. & Cole, G. M. (2005). Curcumin inhibits formation of amyloid β oligomers and fibrils, binds plaques, and reduces amyloid in vivo. *J Biol Chem* **280**, 5892-5901.
- Zacharias, N. & Dougherty, D. A. (2002). Cation- π interactions in ligand recognition and catalysis. *Trends Pharmacol Sci* **23**, 281-287.

References

- Zhang, G. Y., Mills, D. A. & Block, D. E. (2009). Development of chemically defined media supporting high-cell-density growth of *Lactococci*, *Enterococci*, and *Streptococci*. *Appl Environ Microbiol* **75**, 1080-1087.
- Zheng, H., Comeforo, K. & Gao, J. M. (2009). Expanding the fluororous arsenal: tetrafluorinated phenylalanines for protein design. *J Am Chem Soc* **131**, 18-19.
- Zhong, W., Gallivan, J. P., Zhang, Y., Li, L., Lester, H. A. & Dougherty, D. A. (1998). From *ab initio* quantum mechanics to molecular neurobiology: a cation-pi binding site in the nicotinic receptor. *P Natl Acad Sci USA* **95**, 12088-12093.
- Zhou, L., van Heel, A. J. & Kuipers, O. P. (2015). The length of a lantibiotic hinge region has profound influence on antimicrobial activity and host specificity. *Front Microbiol* **6**, 11.

